

20% Multi-Use Poultry Pellet

0163-50

A

Guaranteed Analysis

Crude Protein		(Min)	20.0 %
Crude Fat		(Min)	3.0 %
Crude Fiber		(Max)	5.4 %
Calcium	(Min) 2.2	(Max)	2.7 %
Phosphorus		(Min)	0.7 %
Sodium	(Min) 0.2	(Max)	0.3 %
Methionine		(Min)	0.6 %
Lysine		(Min)	1.0 %
Salt	(Min) 0.3	(Max)	0.3 %
Vitamin A		(Min)	4,156 IU\LB
Vitamin D		(Min)	1,900 IU\LB
Vitamin E		(Min)	24 IU\LB
Niacin		(Min)	36.1 MG
Lactobacillus casei		(Min)	64 mil CFU\LB
Enterococcus faecium		(Min)	64 mil CFU\LB
Bifidobacterium long		(Min)	64 mil CFU\LB
Bacillus licheniform		(Min)	64 mil CFU\LB
Lacto Acidophilus		(Min)	64 mil CFU\LB

Net Weight

50 Lbs (22.7 Kgs)

WP20

List of Ingredients: Corn - Ground, Soybean Meal, Ground Grains, Wheat Millrun, Calcium Carbonate, Rice Bran, Monocalcium Phosphate, Sodium Chloride, DL-Methionine, Choline Chloride, Sodium carboxymethyl-cellulose, and Calcium carbonate., Limestone Ground, Aspergillus Niger Fermentation, Ferrous Sulfate, Zinc proteinate, Manganese Oxide, Selenium Yeast, Zinc Oxide, Manganese Sulfate, Vitamin E Supplement, Niacin Supplement, Vitamin A Supplement, Vitamin D3 Supplement, Calcium Pantothenate, Copper Sulfate, Basic Copper Chloride, Organic Soybean Oil, Riboflavin Supplement, Menadione Nicotinamide Bisulfite, Pyridoxine Hydrochloride (Vitamin B6 Supplement), Organic Rice Hulls, Thiamine Mononitrate, Vitamin B12 Supplement, Ethylenediamine Dihydroiodide, Folic Acid and Biotin, Sodium Bicarbonate, L-Lysine, Niacin, Diatomaceous Earth (flow agent), Hemicellulose Extract, Origanum Oil, Cinnamaldehyde, Thyme White Oil, Hydrolyzed yeast, Dried Distillers Grains, Hydrated Sodium Calcium Aluminosilicate, Dried Aspergillus Oryzae Fermentation Extract, Dried Lactobacillus Acidophilus Fermentation Product, Dried Lactobacillus Casei Fermentation Product, Dried Enterococcus Faecium Fermentation Product, Dried Bifidobacterium Longum Fermentation Product.

A complete feed for layers, broilers, turkeys, ducks, geese, and pheasants older than 8 weeks.

Feeding Directions:

- Feed Free Choice
- Clean feeders and waterers regularly
- Provide appropriate size grit as needed
- Oyster shell or coarse ground limestone should be fed from 16 weeks of age for Laying hens as supplemental calcium.

WARNING: THIS FEED CONTAINS COPPER AT 13 MG/KG. COPPER LEVELS IN THIS FEED MAY BE UNSAFE FOR SHEEP.

Manufactured By: **BAR★ALE** WWW.BARALEINC.COM



Premium Livestock Feeds

1011 Fifth Street,
Williams, CA. 95987
530-473-3333
info@baraleinc.com

