

MEDICATED

0839-50

17% Calf Starter - Medicated

With Decco

For the prevention of coccidiosis caused by Eimeria bovis and Eimeria zuernii.

ACTIVE DRUG INGREDIENT

Decoquinatate 30.4 mg/lb (0.0067 %)

Guaranteed Analysis

Crude Protein	(Min)	17.0 %
Crude Fat	(Min)	2.7 %
Crude Fiber	(Max)	6.6 %
Calcium	(Min)	1.0
Phosphorus	(Min)	0.5 %
Sodium	(Min)	0.6
Potassium	(Min)	1.0 %
Vitamin A	(Min)	12,371 IU\LB
Saccharomyces Cerevi	(Min)	173.98 bil CFU\LB
Lactobacillus casaei	(Min)	174 mil CFU\LB
Enterococcus faecium	(Min)	174 mil CFU\LB
Bifidobacterium long	(Min)	174 mil CFU\LB
Bacillus licheniform	(Min)	174 mil CFU\LB
Lacto Acidophilus	(Min)	174 mil CFU\LB

Net Weight

50 Lbs (22.7 Kgs)

17DEC

List of Ingredients: Soybean Meal, Corn - Rolled, Rolled Barley, Wheat Millrun, Ground Grains, Cane Molasses, Rice Bran, Calcium Carbonate, Sodium Chloride, Organic Wheat Millrun, Organic Grain Screenings, Monocalcium Phosphate, Limestone Ground, Zinc Sulfate, Manganese Oxide, Vitamin E Supplement, Copper Sulfate, Sodium Selenite, Vitamin A Supplement, Vitamin D3 Supplement, Cobalt Sulfate, Calcium Iodate, Zinc proteinate, Manganese proteinate, Copper proteinate, Cobalt proteinate and Calcium carbonate, hydrated calcium potassium sodium aluminosilicates, Sodium Bentonite, Sodium carboxymethyl-cellulose, and Calcium carbonate., Suncured Alfalfa Meal, Diatomaceous earth, Saccharomyces Cerevisae, Selenium Yeast, Hydrolyzed yeast, Brewers Yeast, Hydrated Sodium Calcium Aluminosilicate, Dried Aspergillus Oryzae Fermentation Extract, Dried Lactobacillus Acidophilus Fermentation Product, Dried Lactobacillus Casei Fermentation Product, Dried Enterococcus Faecium Fermentation Product, Dried Bifidobacterium Longum Fermentation Product.

FEEDING INSTRUCTIONS: FEED AT THE RATE OF .7 LBS PER 100 LBS OF BODY WEIGHT TO PROVIDE 22.7 MG

DECOQUINATE PER 100 LBS BODY WEIGHT PER DAY

WARNING: Do not feed to cows producing milk for food. Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates that coccidiosis is likely to be a hazard. Feed type C feed or milk replacer to provide 22.7 mg per 100 lb. bodyweight (0.5 mg/kg) per day.

CAUTION: Do not use in feeds containing bentonite.

CAUTION: DO NOT USE IN FEEDS CONTAINING BENTONITE

This feed contains no more than 0.06% Sodium Bentonite

WARNING: THIS FEED CONTAINS COPPER AT 50 MG/KG. COPPER LEVELS IN THIS FEED MAY BE UNSAFE FOR SHEEP.

WARNING: This feed contains added selenium at 1. mg/kg.

This feed contains selenium over .3PPM Do not feed Selenium at a rate over .3PPM in total ration Do not feed as more than 21.0% of total daily ration.

Do not mix more than 420lbs/Ton in total mixed ration

Manufactured By: **BAR★ALE** WWW.BARALEINC.COM

Premium Livestock Feeds


1011 Fifth Street,

Williams, CA. 95987

530-473-3333

info@baraleinc.com

MEDICATED



8 18314 02066 5